



LJMU Research Online

De Groote, I, Morales, J and Humphrey, L

Oral health in Late Pleistocene and Holocene North West Africa

<http://researchonline.ljmu.ac.uk/id/eprint/8721/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

De Groote, I, Morales, J and Humphrey, L (2018) Oral health in Late Pleistocene and Holocene North West Africa. Journal of Archaeological Science: Reports. ISSN 2352-409X

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Title: Oral Health in Late Pleistocene and Holocene North West Africa

Authors: Isabelle De Groote¹; Jacob Morales², Louise Humphrey³

Affiliations:

1. Liverpool John Moores University, Byrom Street, Liverpool, L3 3AF, UNITED KINGDOM
2. Departamento de Ciencias Históricas, Universidad de Las Palmas de Gran Canaria, Pérez del Toro 1, Las Palmas 35003, SPAIN
3. The Natural History Museum, Cromwell Road, London, SW7 5BD, UNITED KINGDOM

Corresponding author:

Dr Isabelle De Groote
Research Centre in Evolutionary Anthropology and Palaeoecology
School of Natural Sciences and Psychology
Liverpool John Moores University
Byrom Street
Liverpool
L3 3AF
UNITED KINGDOM
Email: i.degroote@ljmu.ac.uk
Office: 0151 2312812

Keywords: Caries, Maghreb, Iberomaurusian, Capsian, diet, subsistence

Highlights:

1. Late Pleistocene populations from Northwest Africa exhibit high caries rates
2. Similarity in poor oral health in Iberomaurusian and Capsian populations
3. Cultural and biological continuity between the Iberomaurusian and Capsian periods.

Funding: This research was supported by the Leverhulme Trust (F/08 735/F) and the Calleva Foundation.

Abstract

Archaeological sites in North West Africa have yielded a rich record of human occupation, including well dated human burials from the Late Pleistocene Iberomaurusian and early Holocene Capsian periods. The transition broadly coincides with climatic amelioration at the end of the Holocene enabling expansion into slighter dryer inland areas. Here we investigate possible changes in oral health and subsistence behaviours during the transition between the Iberomaurusian (n=109 individuals) and the Capsian (n=19 individuals), based on the study of dental remains.

Frequencies in oral pathologies (caries, abscesses, periodontal disease and antemortem tooth loss) were studied to assess possible differences between the late Pleistocene and early Holocene. The Late Pleistocene Iberomaurusians were characterised by high caries frequencies (60% of observed teeth). The Capsians displayed very similar patterns in oral pathologies but a slightly lower percentage of carious teeth (49%).

The similarity in oral health in Iberomaurusian and Capsian populations is consistent with similarities in diet and oral hygiene. The implication of cultural and biological continuity between the Iberomaurusian and Capsian periods is supported by indicators, such as exploitation of wild plants and snails, tooth evulsion and craniofacial and dental morphology.

1. Introduction

1.1 Oral Health

The impact of the adoption of food production as a subsistence pattern on overall health has been the emphasis of a considerable amount of research (Hugot 1968, Turner 1979, Cook 1984, Ruff *et al.* 1984, Bridges 1989, Deacon 1992, Larsen 1995, Jackes *et al.* 1997, Bar-Yosef 1998, Tayles *et al.* 2000, Schollmeyer and Turner 2004, Eshed *et al.* 2006, Starling and Stock 2007, Temple and Larsen 2007, Lukacs 2008, Pinhasi *et al.* 2008, Watson 2008, Kuijt 2009, Pomeroy and Zakrzewski 2009, Eshed *et al.* 2010). This research has been based on the premise that a shift from diverse food resources as part of a hunting and gathering diet to a more restricted agricultural diet consisting predominantly of complex carbohydrates will result in negative health outcomes (Cook 1984, Larsen 1995, Eshed *et al.* 2006, Starling and Stock 2007, Temple and Larsen 2007, Watson 2008, Eshed *et al.* 2010). Increased sedentism in permanent or semi-permanent housing, poor sanitation and the presence of animals as domesticates and pests would have affected the presence and spread of pathogens. Oral pathologies are strongly correlated to subsistence patterns, and have been used by numerous researchers to assess diet, food preparation, and, indirectly, quality of life of past populations (Larsen *et al.* 1991). Teeth are sensitive to fluctuations in diet, pathogenic load, physiological stress, oral hygiene, food preparation, non-dietary use of the teeth, and fluoride intake (Hillson 1996; Larsen 2015). The frequencies of oral pathologies within an osteological sample can provide information about the diet and lifeways of past peoples. Most of these pathological conditions result in the destruction of dental tissues and alveolar bone of both the mandible and maxilla. Changes in oral health and dental wear associated with the transition from a hunter gatherer economy to food production have been documented in the Levant (Eshed *et al.*, 2006), South East Asia (Tayles *et al.*, 2000) Europe (Bennike, 1985; Lubell *et al.*, 1994; Meiklejohn *et al.*, 1998), Japan (Joichi *et al.*, 1996; Oyamada *et al.*, 2004; Temple and Larsen, 2007) and North America (Larsen, 1983; Schollmeyer and Turner, 2004).

Caries, or dental cavities, is the progressive demineralization of dental tissues due to increased acidity resulting from the production of organic acids during fermentation of carbohydrates by bacteria in dental plaque (Hillson 2008). The optimal oral environment for these cariogenic bacteria is achieved by a regular intake of soft, sticky, and sweet foods, in combination with poor oral hygiene (Hillson 1996). The increased reliance on carbohydrate-rich foods, such as wheat and maize, since onset of food production has resulted in an oral environment in which these bacteria can thrive (Hillson 2008; Hillson 1996; Temple 2015). The increased caries frequencies observed during the Neolithic has been linked to the regular consumption of fermentable carbohydrates from cultivated plant foods, such as wheat and other grains. A number of studies on both archaeological and historical samples have demonstrated a relationship between high caries rates and the onset of food production. In general, hunter-gatherers show lower caries rates than those with mixed economy or horticulture (Lanfranco and Eggers 2010). Populations with a greater reliance on carbohydrates, and specifically refined carbohydrates have higher caries rates (Lanfranco and Eggers 2010).

In the Levant, caries frequencies increased slightly from 6.4% in Natufian hunter-gatherers (10,500-8300 BC, non-calibrated) to 6.7% with the onset of the Neolithic (8300-5500 BC) (Eshed *et al.* 2006). In the Americas an increase in caries rates following the transition from horticulture to farming was observed in a number of populations (Cucina *et al.* 2011; Lanfranco and Eggers 2010; Schollmeyer and Turner 2004). Conversely the introduction of rice as a cultivated staple is not always associated with an increase in caries prevalence, reflecting differences in the cariogenicity of carbohydrate staples (Tayles 2009; Tayles *et al.* 2000). At the same time, the importance of carbohydrate-rich wild plant foods by pre-agricultural populations may have been underestimated. Humphrey and colleagues (2014) demonstrated that carbohydrate-rich wild plants such as acorns were widely exploited during the

Late Pleistocene in the Maghreb region. Wild acorns were also commonly consumed by Californian Native Americans and in some groups linked to high caries rates (Turner and Machado 1983).

A series of other dento-alveolar parameters can be recorded to assess oral health in skeletal samples including alveolar resorption caused by periodontal disease, abscesses and antemortem tooth loss. Periodontal disease involves infection the tissues surrounding the tooth and can be either acute or chronic. Acute infections tend to heal and leave few signs after healing. Chronic infections of the gum may result in retraction of the gum line and resorption of the alveolar bone causing exposure of the tooth root above the alveolar margin (Hillson 2001; Nelson 2015). The infection of the alveolar tissue near the apex of the tooth root may result in the accumulation of pus, and is referred to as an abscess. Osteoclastic activity will form a fistula (cavity) inside the bone which may open up to the surface to release pressure through the oral cavity. When there is a clear opening to the fistula in the alveolar bone of a skeleton it is possible to identify this as an abscess (Hillson 2001; Nelson 2015). Any of these infections may lead to antemortem tooth loss. Antemortem tooth loss is identified as a tooth that is missing from the dentition and has a remodelled tooth socket. There are a number of reasons why teeth can be missing including accidental loss, and extensive wear, which results in loosening of the tooth due to continued emergence. In some cases, tooth extraction must be considered a possible cause. Teeth may be extracted to avoid pain and discomfort from an infection, but in late Pleistocene and early Holocene Maghreb teeth were often deliberately removed for cultural reasons. The deliberate removal of healthy teeth is referred to as evulsion or ablation. The majority of individuals from the sites considered here show the deliberate removal of two or more incisors (De Groote and Humphrey 2015; De Groote and Humphrey, 2017; Humphrey and Bocaegé 2008). The widespread frequency of the practice of evulsion across the region indicates that the absence of anterior teeth in these samples is likely due to cultural tradition, rather than an oral pathology.

1.2 Archaeological context

Research on the transition from the Late Pleistocene (Iberomaurusian) to early Holocene (Capsian) in the Maghreb has focused on changes in lithic assemblages (Linstädter 2008; Linstädter 2011; Lubell et al. 1994; Lubell et al. 1984) and variation in skeletal and dental morphology (Irish 2000; Lubell et al. 1994; Lubell et al. 1984; Sheppard and Lubell 1991) but the transition has not been considered in relation to human diet and health. The study of oral health can provide an additional insight into the nature of this transition since the type and prevalence of oral pathologies are directly influenced by dietary intake and food processing (Hillson 2001; Hillson 1996; Larsen et al. 1991; Lukacs 2008).

Recent research highlighted a very high prevalence of caries in Late Pleistocene hunter –gatherers from the Iberomaurusian site of Grotte des Pigeons, Taforalt, Morocco (Humphrey et al. 2014). More than fifty-one percent of all observed adult teeth had at least one carious lesion. Charred macrobotanical remains from occupational deposits revealed evidence for the use of numerous wild plant foods with a preference for large-sized and potentially stored seeds (Morales, this volume). The most abundant macrobotanical remains found in the site were the fragments of acorn (*Quercus ilex*), pine nuts (*Pinus pinaster*) and juniper (*Juniperus phoeniceae*). Frequent consumption of wild plant foods such as acorns that are rich in fermentable carbohydrates would have contributed to the poor oral health observed in the dental series from Taforalt (Humphrey et al. 2014). This study will expand upon the previously published results by ascertaining whether the patterns of oral pathology (prevalence and distributions of carious lesions, antemortem tooth loss, abscesses, and alveolar resorption) observed at Taforalt were typical for the Iberomaurusian period in the Maghreb. Secondly, we investigate whether the transition from the Late Pleistocene Iberomaurusian to the early Holocene Capsian brought with it a change in oral health and subsistence behaviours.

The Iberomaurusian is a Late Pleistocene Later Stone Age industry dominated by small backed blades and backed points (Barton et al. 2013). The term is also used for the people who made this industry. Recent re-analyses of the lithic technology and a comprehensive re-dating programme and evaluation of existing dates, have dated the earliest occurrence of the Iberomaurusian industry at ~25 ka cal BP and most recent occurrence at ~11.5ka cal BP (Hogué and

Barton 2016). Within the Iberomaurusian stone tool industry two subdivisions can be identified with the older one recognised by Ouchtata retouched blades, and a more recent one dominated by microlithic backed bladelets (Barich and Garcea 2008; Barton et al. 2013). Microlithic components were present at Grotte des Pigeons, Taforalt by 15,500 – 15,000 cal BP (Barton et al. 2013; Hogue and Barton 2016).

At Grotte des Pigeons, Taforalt, (hereafter referred to as Taforalt) the dominant presence of microlithic components occurred shortly before an abrupt sedimentary transition. The overlying Grey Series deposits are characterised by the rapid accumulation of an ashy midden, containing vast quantities of ash, burnt bone, snail shells and lithic debitage, indicating an intensification of human activity around ~15 ka cal BP (Barton et al. 2013). The ash-rich deposits also preserve abundant evidence for harvesting and consumption of wild plant foods such as acorns, pine nuts and juniper berries, which supplemented the animal protein recovered from medium-sized mammals and molluscs (Humphrey et al. 2014; Taylor et al. 2011). The presence of charred rhizomes from Alfa grass/esparto (*Stipa tenacissima*) in the levels investigated is indirect evidence that the Iberomaurusians were producing baskets and other woven pieces of equipment that could have been used to collect wild food items (Morales, this volume). Grindstones have also been recovered from the burial deposits (Humphrey et al. 2014; Roche 1963). Similar changes in sedimentation, suggesting intensification of human occupation during the Iberomaurusian, occur at other sites in Morocco, including Ifri n’Ammar and Ifi El-Baroud (Linstädter et al. 2012).

The Capsian is an early Holocene stone tool industry found primarily at sites in the Eastern Maghreb (modern day Tunisia and Algeria) between ~ 11.5 and 7.0 ka cal BP (Linstädter 2008). The term also refers the people who occupied those sites. The Capsian lithic industry is based on the production of bladelets but distinguished from the Iberomaurusian by the inclusion of small geometric microliths (Lubell 2001). Two typologically distinct industries have been identified: The Typical and the Upper Capsian (Jackes and Lubell 2008; Lubell 2001; Lubell et al. 1976; Lubell et al. 1984; Merzoug 2014; Rahmani 2004). Both have a tool assemblage consisting of backed blades, scrapers and burins, as well as some bone tools. The subdivisions were initially interpreted as consecutive, but are now generally recognised as contemporaneous (Linstädter 2008; Rahmani 2004). The Upper Capsian is characterised by increased variability in raw materials and more efficient use of prepared tools (using a pressure detachment technique (Mulazzani et al.)), and may reflect the movement of Capsian people into more remote regions where high quality raw materials were less abundant (Linstädter 2008; Rahmani 2004). Capsian open air sites, often referred to as *escargotières*, are characterised by mounds of accumulated ash, flint tools, burnt bone and stone and vast quantities of land snail shells. Evidence for harvesting of wild food resources including land snails and plants and hunting of wild animals points to a hunter-gatherer lifestyle (Morales, this volume, Aouadi et al. 2014).

2. Materials and Methods

2.1 Materials

The skeletal and dental sample that was used for this study was assembled from a number of Iberomaurusian and Capsian sites (Figure 1). The number of individuals, number of analysed teeth and sample breakdown into age categories is presented in Table 1.

>> Figure 1 and Table 1

The Iberomaurusian sample comes from three well-dated Late Pleistocene sites in Morocco and Algeria. Historic and recent excavations at Grotte des Pigeons in Taforalt in Morocco revealed numerous partially articulated human skeletons and disarticulated skeletal parts from primary, secondary and disturbed burials in a spatially demarcated area towards the back of the cave (Balout 1954; Ferembach 1962; Humphrey et al. 2012; Mariotti et al. 2009;

Mariotti et al. 2014). Direct dates on human bone from the recent phase of excavations have yielded age estimations between 15,077 cal BP and 13,892 cal BP (Humphrey et al. 2014), but burials may have continued throughout the Iberomaurusian period. The most recent Iberomaurusian deposits at Grotte des Pigeons are dated to around 12,600 cal BP (Barton et al. 2013). The second Iberomaurusian sample is from the rock shelter of Afalou Bou Rhummel in Algeria (hereafter referred to as Afalou) (Arambourg et al. 1934; Hachi 1996). The samples studied here were recovered during excavations between 1928 and 1929 (Arambourg et al. 1934). The majority of individuals were found disarticulated and incomplete in Level 1. More recently partially disarticulated human skeletons representing approximately eight individuals were found in a separate burial chamber and these yielded dates of 13,120 +/- 370 BP (Alger 0008), 12,400 +/- 240 BP (Ly 3228) and 12,020 +/- 170 BP (Gif 6532) (Hachi 1996). A single individual from Dar-es-Soltan was also included with a date of 13,400 +/- 700 YA (OSL4-X2402) (Schwenninger et al. 2010).

The Capsian sample is smaller than the Iberomaurusian sample and comprises individuals from nine Typical and Upper Capsian sites in Eastern Algeria and Tunisia (Table 1). Site 12 (Ain Berriche) is an escargotière in Algeria with a Capsian industry (7753-8539 cal BP (Jackes and Lubell 2014)). Six adult and two juvenile skeletons were excavated from the Minnesota trench during excavations led by Jenks in 1930 (Haverkort and Lubell 1999). One of the adult skeletons was considered intrusive at the time of excavation and this has recently been confirmed by radiocarbon dating (Jackes et al 2015). Two other adult skeletons yielded calibrated radiocarbon dates between 9000 and 8700 cal BP (Jackes et al 2015). Dentitions from three of the adult skeletons and an additional fragment were included in this study. Ain Dokkhara is an escargotière in Algeria with an Upper Capsian industry. Excavations in 1949 revealed a well preserved skeleton of a young male in deposits that has been indirectly dated at 6580 +/- 100 BC, based on a radiocarbon date on land snails from the same horizon (Chamla 1973). The skeleton of an adult male aged between 40 and 50 years was found during excavations at Ain Meterchem an escargotière in Tunisia in 1948. Some authors have considered this to be an intrusive burial dating to the same Neolithic stage as other bead decorated burials (Camps 1974, but Chamla 1978), but Vallois and Felice (1979) have made a strong case that the skeleton belongs to the Capsian. Disarticulated human bones from several individuals were uncovered during excavations at Grotte des Hyènes between 1925 and 1926, including one relatively complete skull that had undergone evulsion of all eight incisors (Royer 1926, Arambourg et al. 1934, Balout 1954) and upper or lower jaws from several other individuals. A fragmentary female skull was found at Koudiat-el-Kherrouba, an escargotière in Algeria during excavations in 1933 (Arambourg et al. 1934, Balout 1954). Some authors have suggested that the context may be Neolithic (Briggs 1955) but the skull shows evidence of evulsion of all eight incisors which is typical for the Capsian period (Humphrey and Bocaeg 2008). Several skulls were found at Khanguet el Mouhaâd, an escargotière in Algeria, but only one of these, is a young adult female found in 194, is considered to be in a secure Upper Capsian context (Briggs 1953). Mechta el Arbi is an escargotière in Algeria with an Upper Capsian industry. A large number of human remains were recovered in a series of excavations between 1907 and 1927 (Balout 1954) including four individuals included in this study. Human bones found at Mesloug, an escargotière in Algeria, in 1928 (Balout 1954) included a partial mandible with evulsion of the lower central incisors that was included in this study. Despite some uncertainty over its cultural association, a partial skull from Redeyef, a rock shelter in Tunisia first reported in 1912 (Balout 1954) was included in this study.

Preservation was variable across the sample, with some individuals represented by complete skeletons and others only by fragmentary dental remains. Fragmentary dentitions were evaluated for caries prevalence and distribution to maximise sample size. The sample was divided into four age categories (see below). It was not possible to make comparisons between males and females because there were too few reliably sexed dentitions. No statistical analyses were undertaken between time periods due to the small and uneven sample sizes.

2.2 Methods

2.2.1 Age estimation

Pubic symphysis morphology, auricular surface morphology, ectocranial suture closure, and sternal rib end changes were used to assign individual skeletons into one of three age categories: young adult (18-25), middle adult (25-55) and old adult (55+) (Buikstra & Ubelaker, 1994). Those dentitions unable to be assigned an age category were grouped into an undetermined age category. Juveniles were excluded from the study. For samples without associated skeletal material, age estimation was carried out using dental wear stages.

2.2.2 Dento-alveolar parameters

A series of dento-alveolar parameters were recorded by the same observer (IDG): caries, antemortem tooth loss (AMTL), postmortem tooth loss (PMTL), abscesses, alveolar bone resorption, and tooth wear. All teeth were examined with good lighting and a magnification of 4x and 10x. The frequency of each pathology is reported as a proportion of its observable parameter (e.g. number of sockets). The number of observations for each of these observable parameters depends on the presence and preservation of teeth and alveoli and which pathology is considered.

- The *observable sockets* are the combination of all teeth and alveoli observed in the sample.
- The *valid observed sockets* are the observable sockets minus sockets affected by post-mortem tooth loss.
- The *observed teeth* are the observable sockets minus sockets affected by post-mortem and ante-mortem tooth loss.

Alveolar resorption

Alveolar bone resorption was recorded as present if at least a 2mm distance was measured between the crest of the alveolar bone and the cemento-enamel junction on any surface of the tooth crown (mesial, distal, lingual or buccal) (Hillson, 1996). Root exposure is reported as a proportion of observed teeth.

Postmortem tooth loss (PMTL)

Postmortem tooth loss was defined as an empty socket showing no sign of alveolar remodelling (Nelson 2015; Ortner and Putschar 1981). PMTL is reported as the proportion of observable sockets.

Caries

For these analyses a carious lesion was only considered present when there was clear penetration and tissue breakdown of the enamel (Hillson 2001). The number of lesions and locations was scored for each observed tooth. For overall caries frequency, only one surface (occlusal, mesial, distal, lingual or buccal) of the tooth had to be affected for caries to be considered present on the tooth. Caries are expressed as the proportion of observed teeth.

Caries categories were expressed as the proportion of carious teeth.

- Gross caries was determined present when a lesion affected more than one tooth surface and the surface of origin of the caries could no longer be determined.
- Crown caries refer to the presence of caries on one of more of these locations on the crown: (a) occlusal molar fissures, (b) occlusal attrition caries in the dentine, (c) buccal surface, (d) lingual surface, (e) mesial surface, (f) distal surface, (g) pit caries, and (h) rim chipping caries.
- Root caries refer to the presence of at least one carious lesion on one or more of these locations on the root: (a) buccal surface, (b) lingual surface, (c) mesial surface, (d) distal surface.

Antemortem tooth loss (AMTL)

Antemortem tooth loss was defined as an empty socket showing any sign of alveolar remodelling (Nelson 2015; Ortner and Putschar 1981). Because of the high prevalence of evulsion of the anterior teeth in these populations, we report AMTL only as post-canine AMTL. AMTL is reported as the proportion of valid observed post-cranial sockets.

Abscesses

The distinction between abscesses, granulomas and cysts is difficult to make in an archaeological assemblage without x-ray observations (Hillson 1996; Nelson 2015). Therefore all periapical cavities are grouped here under the term abscess. Abscesses were recorded by direct observation only and defined as a periapical cavity with relatively smooth walls. Because of the lack of X-ray observations for these cavities their prevalence will be underestimated. We focused on periapical bone loss that was visible by the naked eye only. Abscesses were recorded as present when there was a clear cavity in the alveolar bone and reported as the proportion of observable sockets.

Alveolar bone resorption (root exposure)

Periodontal disease often results in bone resorption and root exposure. It was recorded as present if at least a 2mm distance was measured between the crest of the alveolar bone and the cemento-enamel junction on any tooth surface (mesial, distal, lingual or buccal). Root exposure is reported as the proportion of observed teeth exhibiting alveolar bone resorption.

3. Results

In total 128 individual dentitions from the Iberomaurusian (N=109) and Capsian (N=19) were represented by at least one tooth or alveoli (Table 2 and 3). The number of observable tooth sockets for the Iberomaurusian was 1991 and for the Capsian 404. A summary of the number of individuals and observed sockets and teeth is presented in Table 2 and Table 3 together with frequencies of PMTL for the postcanine dentition. PMTL affected 334 (17%) of Iberomaurusian and 47 (12%) of Capsian observable sockets. Due to the near-universal practice of tooth evulsion in this sample AMTL is only presented for the post-canine dentition. The total number of post-canine sockets observed with AMTL was 112 (9% of valid observable sockets) for the Iberomaurusian (35% of individuals) and 34 (14% of valid observable sockets) for the Capsian (50% of individuals). The total number of surviving observable teeth was 1417 for the Iberomaurusian and 271 for the Capsian.

>> Table 2

>> Table 3

Comparisons between the 951 observable sockets from Taforalt and 1024 observable sockets from Afalou (Table 4) demonstrates that the high caries frequencies observed for Grotte des Pigeon (51%, n=345/681 observed teeth) reported by Humphrey et al. (2014) are actually surpassed by Afalou (69%, n=498/727 observed teeth). Caries rates for the all age groups from Afalou were higher than at Taforalt (50% compared to 33% of observed teeth respectively for young adults, 73% compared to 55% for middle adults and 80% vs 53% for old adults). These results suggest that high caries frequencies may have been characteristic of the time-period represented by these samples.

>> Table 4

The caries frequencies for the Iberomaurusians and Capsians are reported in Table 2 and 3. Both samples presented very high caries frequencies with 60% (n=844/1417) of observed teeth affected in the Iberomaurusian and 49% (133/271) of observed teeth affected in the Capsian. In total, 97% of the 109 Iberomaurusian dentitions and 74% of 19 Capsian dentitions presented at least one carious lesion. Crown caries were the most frequent with more than

30% of observed teeth affected in both time periods. The frequency of root caries was similar during the Iberomaurusian (38% of observed teeth) and Capsian (35% of observed teeth).

Sample sizes for the Capsian are small so comparison across the age categories may be affected by sample bias. Nevertheless it is evident that caries affected young and old adults and that even gross caries were observed amongst all age groups (Table 2). In the Iberomaurusian sample the presence of gross caries followed the expected pattern, increasing between age groups. Post-canine ante-mortem tooth loss followed the same pattern suggesting that caries was an underlying cause.

The prevalence of non-carious oral pathologies is presented in Table 2. Alveolar resorption was present 51% of observed Iberomaurusian teeth and was common even amongst young adults. A similar proportion of Capsian teeth (58%) were affected and, as in the Iberomaurusian, all age categories were affected. Abscesses were present in 37% of Iberomaurusian dentitions and 63% of Capsian dentitions, representing 4% and 7% of observable sockets respectively.

3. Discussion

This research sought to determine whether there are differences in oral pathology between the Iberomaurusian and the Capsian periods and whether the high caries frequencies previously reported for the dental series from Grotte des Pigeons at Taforalt are unique to this site, or whether they are representative of that time period. Comparisons of the dental series from Taforalt and Afalou (Algeria), shows that caries rates are slightly higher at Afalou than at Taforalt. Comparisons between the Iberomaurusian and Capsian samples show that caries rates were slightly lower in the Capsian (59.6% (IB) vs 49.1% (CAP)), but were still very high compared to published caries frequencies for other hunter-gatherers populations (Figure 2) (Lanfranco and Eggers 2010).

>> Figure 2

Reported caries frequencies will be inflated in these samples due to the absence of anterior teeth caused by the widespread practice of evulsion during the Iberomaurusian and Capsian (Humphrey and Bocaage 2008, De Groote and Humphrey 2017). Clinical research has demonstrated that mandibular central incisors are the least likely to be carious, followed by canines and premolars, while molars were the most likely affected by caries (Demirci et al 2010). This observation must be kept in mind when comparing these results to those reported for populations that do not practice evulsion.

Caries causes are multifactorial and complex. The composition and the proportion of carbohydrates consumed by the individual plays a major role. Other factors that contribute to differences in oral health between populations are tooth crown morphology, fluoride content of the local groundwater, and the composition of oral microbiota, oral hygiene and transmission of cariogenic bacteria within the population. Variation in tooth morphology, in particular the number and patterning of cusps and degree of crenulation, can have a major impact on caries susceptibility (Hillson 2001). Detailed analysis of dental morphology has shown that Iberomaurusian and Capsian populations exhibit similar and simple tooth morphology (Irish 2000) such that the effect of tooth morphology on caries development is likely to be equivalent across all samples. Fluoride content in the groundwater in the Maghreb is unlikely to have varied over time, but may show local variation. Studies show there is an excess of fluoride in the groundwater, which has resulted in fluorosis being an endemic disease in this region (Brunt et al. 2004; Fantong et al. 2010) but there was no evidence of fluorosis in the Iberomaurusian and Capsian dentitions.

Poor oral hygiene can lead to a more rapid progression of tooth decay. A lack of tooth cleaning to remove the built-up plaque would have contributed to the detrimental effect of these sticky carbohydrates). Tayles et al (2000) concluded that frequent snacking was the cause of higher caries rates in females compared to males despite

consuming the same diet (Tayles et al. 2000). Regular snacking on carbohydrate rich plants or derived foods such as breads and porridges, may have contributed to the extremely high number of oral pathologies in this group.

Caries does not exist without the presence and transmission of oral caries-causing microbiota such as *Streptococcus mutans*. Adler and colleagues (Adler et al. 2013) analysed oral bacteria in dental calculus from archaeological assemblages and concluded that the prevalence and virulence of cariogenic oral bacteria increased during the Neolithic. The evidence presented here for high caries frequencies in hunter-gatherer Late Pleistocene and early Holocene populations from the Maghreb suggests that the origin of highly virulent microbiota predated the Neolithic in North West Africa. Close contact between individuals would have facilitated both horizontal and vertical transmission of cariogenic bacteria (Humphrey et al. 2014). These results warrant the revision of the origins of caries as an epidemic.

The similarity in oral health between Iberomaurusian and Capsian populations implies similarities in subsistence behaviour and oral hygiene and indicates the presence of cariogenic bacterial in the oral microbiota. These similarities are consistent with cultural and biological continuity between the Iberomaurusian and Capsian periods. Charred macro-botanical remains recently found in sites from both periods indicate that carbohydrate-rich plants were extensively processed for human subsistence, and thus confirm that consumption of those could account for the high caries attested (Humphrey et al 2014; Morales, this volume). Acorns, pine nuts, juniper berries, wild pistachios, wild legumes and others, up to 20 different taxa, appear abundantly in layers dated after ~15 ka cal BP. Preservation conditions of the vegetal remains suggest that heat processing and storage could be carried out, increasing the energetic value of plant foods. The presence of wild plants is also recorded in Neolithic deposits from those sites, indicating that, despite the appearance of farming, wild plants kept a very important role in the diet of this period (Mulazzani et al 2016; Morales, this volume). Evidence for systematic exploitation of land molluscs at late Late Pleistocene and early Holocene sites in both Eastern and Western Maghreb is a further indication of cultural similarity in diet between the Iberomaurusian and Capsian. Iberomaurusian deposits at Taforalt show evidence for harvesting of five species of edible snails (Taylor et al. 2011). Shells of terrestrial gastropods are also recorded in epipaleolithic and Neolithic levels at numerous other cave sites in North East Morocco (Hutterer et al 2014). Exploitation of edible land mollusc has long been considered characteristic of the Capsian period, to the extent that Capsian open-air sites are often known as “escargotieres”, in reference to vast quantities of snail shells recovered from these sites (Aouadi et al. 2014; Hutterer et al. 2014; Lubell 2004a; Lubell 2004b).

Cultural indicators also support similarities between the Iberomaurusian and Capsian. The technocomplex of both periods is closely related, involving the use of single platform cores for bladelet production (Sheppard and Lubell 1991). The practice of tooth evulsion, or the deliberate removal of healthy teeth, is also characteristic of both periods. During the Iberomaurusian period, most individuals practiced evulsion of both upper central incisors. In the Capsian, this practice became more extreme and in some cases all anterior teeth were removed (De Groote and Humphrey 2015; De Groote and Humphrey 2017; Humphrey and Bocaeye 2008). Recent studies of dental morphology are also consistent with a degree of long-term biological similarity in the Maghreb (Irish 2000). Iberomaurusian dentitions already exhibit many of the morphological features present in the North African Dental Trait Complex and share these features with all more recent populations from North West Africa (Irish 2000).

4. Conclusion

This paper set out to investigate whether the transition from the Late Pleistocene Iberomaurusian to the early Holocene Capsian from the Maghreb region brought with it a change in oral health and subsistence behaviours. The results showed high oral pathologies in both populations exceeding those observed in many food producing populations, although those in the Capsian were somewhat lower than those of the Iberomaurusian. The increased reliance on carbohydrates with the onset of food production and their detrimental effect on oral health are well documented in both archaeological and historical samples but both phenomena appear to have predated food production in the Maghreb. The contribution of wild plant foods contributed to Iberomaurusian and Capsian diets is

explored elsewhere in this volume (Morales, this volume). The results also suggest that the origin of highly virulent microbiota predated the onset of the Neolithic in North West Africa.

To conclude, the subsistence pattern of the Late Pleistocene Iberomaurusians of North West Africa incorporated a significant component of wild plant foods which was associated with high frequencies of oral pathologies. Early Holocene Capsian populations exhibit similarities in oral pathologies. Other cultural indicators, such as a closely related technocomplex, the exploitation of molluscs and the widespread practice of dental evulsion, suggest further parallels between the time periods.

Acknowledgements

For access to collections we thank David Lubell and Mary Jackes, William Green and Nicolette Meister from the Logan Museum at Beloit College, Professor de Lumley and Amélie Vialet from l'Institut de Paléontologie Humaine, Kieran McNulty at the University of Minnesota and Abdeljalil Bouzouggar from the National Institute of Archaeological Science and Heritage (INSAP, Morocco). We are grateful to Abdeljalil Bouzouggar, Nick Barton, Ali Freyne, Peter Berridge and Paul Berridge for assistance and discussions.

REFERENCES

- Adler CJ, Dobney K, Weyrich LS, Kaidonis J, Walker AW, Haak W, Bradshaw CJ, Townsend G, Sołtysiak A, and Alt KW. 2013. Sequencing ancient calcified dental plaque shows changes in oral microbiota with dietary shifts of the Neolithic and Industrial revolutions. *Nature genetics* 45(4):450-455.
- Aouadi N, Dridi Y, and Ben Dhia W. 2014. Holocene environment and subsistence patterns from Capsian and Neolithic sites in Tunisia. *Quaternary International* 320(0):3-14.
- Arambourg C, Boule M, Vallois HV, and Verneau R. 1934. Les grottes paléolithiques des Beni Segoual, Algérie: Masson.
- Balout L. 1954. Les Hommes Préhistoriques du Maghreb et du Sahara. *Libyca* 2:214-424.
- Bar-Yosef O. 1998. The Natufian culture in the Levant, threshold to the origins of agriculture. *Evolutionary Anthropology* 6(5): 159-177.
- Barich B, and Garcea EA. 2008. Ecological Patterns in the Upper Pleistocene and Holocene in the Jebel Gharbi, Northern Libya: Chronology, Climate and Human Occupation. *African Archaeological Review* 25(1-2):87-97.
- Barton RNE, Bouzouggar A, Hogue JT, Lee S, Collcutt SN, and Ditchfield P. 2013. Origins of the Iberomaurusian in NW Africa: New AMS radiocarbon dating of the Middle and Later Stone Age deposits at Taforalt Cave, Morocco. *Journal of Human Evolution* 65(3):266-281.
- Bennike P. 1985. *Palaeopathology of Danish skeletons: a comparative study of demography, disease and injury*: Akademisk Forlag.
- Bridges PS. 1989. Changes in Activities with the Shift to Agriculture in the Southeastern United States. *Current Anthropology*. 30(3): 385.
- Briggs, L.C. 1953. Tête osseuse du Khanguet-el-Mouhaâd. *Libyca* 1: 120-141.
- Briggs LC. 1955. The stone age races of Northwest Africa. *Bulletin of the American School of Prehistoric Research* (18):1-98.
- Brunt R, Vasak L, and Griffioen J. 2004. Fluoride in groundwater: Probability of occurrence of excessive concentration on global scale. Utrecht: International Groundwater Resources Assessment Centre. p 1-12.
- Buikstra JE and Ubelaker DH. 1994. Standards for data collection from human skeletal remains.
- Camps G. 1974. Les civilisations préhistoriques de l'Afrique du Nord et du Sahara. Paris: Doin. 336 p.
- Chamla M-C. 1973. Etude anthropologie de l'homme capsien de l'Aïn Dokkara (Algérie orientale). *Libyca* 21:9-55.
- Chamla MC. 1978. Le peuplement de l'Afrique du Nord de l'épipaléolithique B l'époque actuelle. *L'Anthropologie* 82:385-430.
- Cook DC. 1984. Subsistence and health in the Lower Illinois Valley: osteological evidence' in *Paleopathology at the origins of agriculture*. Academic Press, 235-269.
- Cucina A, Cantillo CP, Sosa TS, and Tiesler V. 2011. Carious lesions and maize consumption among the Prehispanic Maya: An analysis of a coastal community in northern Yucatan. *American Journal of Physical Anthropology* 145(4):560-567.
- De Groote I and Humphrey LT. 2015. Characterizing evulsion in the Later Stone Age Maghreb: Age, sex and effects on mastication. *Quaternary International*.
- De Groote I and Humphrey LT. 2017. The chronology and significance of ablation in the Later Stone Age Maghreb. In : Eds. Burnett, S.E. and Irish, J.D. *A World View of Bioculturally Modified Teeth*. Gainesville, FL. University Press of Florida. p 19-32.
- Deacon HJ. 1992. Southern Africa and Modern Human Origins. *Philosophical Transactions: Biological Sciences*, 337(1280): 177-183.
- Demirci, M., Tuncer, S. and Yuceokur, A.A., 2010. Prevalence of caries on individual tooth surfaces and its distribution by age and gender in university clinic patients. *European journal of dentistry*, 4(3), p.270.
- Eshed V, Gopher A, and HersHKovitz I. 2006. Tooth wear and dental pathology at the advent of agriculture: New evidence from the Levant. *American Journal of Physical Anthropology* 130(2):145-159.
- Eshed, V., Gopher, A. and HersHKovitz, I. 2006. Tooth wear and dental pathology at the advent of agriculture: New evidence from the Levant. *American Journal of Physical Anthropology*, 130(2), 145-159.
- Eshed, V., Gopher, A., Pinhasi, R. and HersHKovitz, I. 2010. Paleopathology and the origin of agriculture in the Levant. *American Journal of Physical Anthropology*, 143(1), 121-133.
- Fantong W, Satake H, Ayonghe S, Suh E, Adelana S, Fantong E, Banseka H, Gwanfogbe C, Woincham L, Uehara Y et al. 2010. Geochemical provenance and spatial distribution of fluoride in groundwater of Mayo Tsanaga River

- Basin, Far North Region, Cameroon: implications for incidence of fluorosis and optimal consumption dose. *Environmental Geochemistry and Health* 32(2):147-163.
- Ferembach D. 1962. La Nécropole Epipaléolithique de Taforalt (Maroc Oriental): Étude des squelettes humains. Edita Casablanca, Rabat. 171 p.
- Hachi S. 1996. L'Ibéromaurusien, découverte des fouilles d'Afalou (Bédjaïa, Algérie). *L' Anthropologie* 100(1):55-76.
- Haverkort CM, and Lubell D. 1999. Cutmarks on Capsian human remains: Implications for Maghreb Holocene social organization and palaeoeconomy. *International Journal of Osteoarchaeology* 9(3):147-169.
- Hillson S. 2001. Recording dental caries in archaeological human remains. *International Journal of Osteoarchaeology* 11(4):249-289.
- Hillson S. 2008. The current state of dental decay. In: Irish JD, and Nelson G, editors. *Technique and application in dental anthropology*. Cambridge: Cambridge University Press. p 111-136.
- Hillson SW. 1996. *Dental Anthropology*. Cambridge Cambridge University Press.
- Hogue J, and Barton R. 2016. New radiocarbon dates for the earliest Later Stone Age microlithic technology in Northwest Africa. *Quaternary International*.
- Hugot H J. 1968 The Origins of Agriculture: Sahara. *Current Anthropology*, 9(5): 483-488.
- Humphrey L, Bello SM, Turner E, Bouzouggar A, and Barton N. 2012. Iberomaurusian funerary behaviour: Evidence from Grotte des Pigeons, Taforalt, Morocco. *Journal of Human Evolution* 62(2):261-273.
- Humphrey L, and Bocaëge E. 2008. Tooth Evulsion in the Maghreb: Chronological and Geographical Patterns. *African Archaeological Review* 25(1):109-123.
- Humphrey LT, De Groote I, Morales J, Barton N, Collcutt S, Bronk Ramsey C, and Bouzouggar A. 2014. Earliest evidence for caries and exploitation of starchy plant foods in Pleistocene hunter-gatherers from Morocco. *Proceedings of the National Academy of Sciences* 111(3):954-959.
- Hutterer R, Linstädter J, Eiwanger J, and Mikdad A. 2014. Human manipulation of terrestrial gastropods in Neolithic culture groups of NE Morocco. *Quaternary International* 320(0):83-91.
- Irish JD. 2000. The Iberomaurusian enigma: North African progenitor or dead end? *Journal of Human Evolution* 39(4):393-410.
- Jackes M, and Lubell D. 2008. Early and Middle Holocene Environments and Capsian Cultural Change: Evidence from the Télijdjène Basin, Eastern Algeria. *African Archaeological Review* 25(1):41-55.
- Jackes M, and Lubell D. 2014. Capsian mortuary practices at Site 12 (Aïn Berriche), Aïn Beïda region, eastern Algeria. *Quaternary International* 320:92-108.
- Jackes M, Lubell D. and Crann, C. 2015. Dating and redating Capsian skeletons 3A-4 and 3A-7, Aïn Berriche (Algeria) *Journal of Archaeological Science: Reports* 4 (2015) 78–85.
- Jackes, M., Lubell, D. and Meiklejohn, C. 1997. Healthy but mortal: human biology and the first farmers of western Europe. *Antiquity* 71(273): 639-658.
- Joichi O, Yoshitaka M, Yoshikazu K, and Atsushi R. 1996. Dental Morbid Condition of Hunter-Gatherers on Okinawa Island during the Middle Period of the Prehistoric Shell Midden Culture and of Agriculturalists in Northern Kyushu during the Yayoi Period. *Anthropological science : journal of the Anthropological Society of Nippon* 104(4):261-280.
- Kuijt I. 2009. What Do We Really Know about Food Storage, Surplus, and Feasting in Preagricultural Communities?, *Current Anthropology*, 50(5), 641-644.
- Lanfranco LP, and Eggers S. 2010. The usefulness of caries frequency, depth, and location in determining cariogenicity and past subsistence: A test on early and later agriculturalists from the Peruvian coast. *American Journal of Physical Anthropology* 143(1):75-91.
- Larsen CS. 1983. Behavioural implications of temporal change in cariogenesis. *Journal of Archaeological Science* 10(1):1-8.
- Larsen CS. 1995. Biological Changes in Human Populations with Agriculture. *Annual Review of Anthropology* 24:185-213.
- Larsen, C.S., 2015. *Bioarchaeology: interpreting behavior from the human skeleton* Cambridge University Press.
- Larsen CS, Shavit R, and Griffin MC. 1991. Dental Caries Evidence for Dietary Change: An Archaeological Context. In: Kelley MA, and Larsen CS, editors. *Advances in Dental Anthropology*. New York: Wiley-Liss. p 179-202.
- Linstädter J. 2008. The Epipalaeolithic-Neolithic-Transition in the Mediterranean region of Northwest-Africa. *Quartär* 55:41-62.

- Linstädter J. 2011. The Epipalaeolithic-Neolithic Transition in the Eastern Rif Mountains and the Lower Moulouya valley, Morocco. The Last Hunter-Gatherers and the First Farming Communities in the South of the Iberian Peninsula and North of Morocco Proceedings of the Work Shop Faro 2:e4.
- Linstädter J, Eiwanger J, Mikdad A, and Weniger GC. 2012. Human occupation of Northwest Africa: A review of Middle Palaeolithic to Epipalaeolithic sites in Morocco. *Quaternary International* 274:158-174.
- Lubell D. 2001. Late Pleistocene-Early Holocene Maghreb. In: Peregrine PN, and Ember M, editors. *Encyclopedia of Prehistory*. New York: Kluwer Academic/Plenum Publishers.
- Lubell D. 2004a. Are land snails a signature for the Mesolithic-Neolithic transition. *Documenta Praehistorica* 31(1):1-24.
- Lubell D. 2004b. Prehistoric edible land snails in the circum-Mediterranean: the archaeological evidence. *Petits animaux sociétés humaines: du complément alimentaire aux ressources utilitaires*:77-98.
- Lubell D, Hassan FA, Gautier A, and Ballais J-L. 1976. The Capsian Escargotieres. *Science* 191(4230):910-920.
- Lubell D, Jackes M, Schwarcz H, Knyf M, and Meiklejohn C. 1994. The Mesolithic-Neolithic Transition in Portugal: Isotopic and Dental Evidence of Diet. *Journal of Archaeological Science* 21(2):201-216.
- Lubell D, Sheppard P, and Jackes M. 1984. Continuity in the Epipalaeolithic of northern Africa with an emphasis on the Maghreb. In: Wendorf F, and Close A, editors. *Advances in World Archaeology*. New York: Academic Press.
- Lukacs John R. 2008. Fertility and Agriculture Accentuate Sex Differences in Dental Caries Rates. *Current Anthropology* 49(5):901-914.
- Mariotti V, Bonfiglioli B, Facchini F, Condemi S, and Belcastro MG. 2009. Funerary practices of the Iberomaurusian population of Taforalt (Tafoughalt; Morocco, 11-12,000 BP): new hypotheses based on a grave by grave skeletal inventory and evidence of deliberate human modification of the remains. *Journal of Human Evolution* 56(4):340-354.
- Mariotti V, Condemi S, and Belcastro MG. 2014. Iberomaurusian funerary customs: new evidence from unpublished records of the 1950s excavations of the Taforalt necropolis (Morocco). *Journal of Archaeological Science* 49:488-499.
- Meiklejohn C, Petersen EB, and Alexandersen V. 1998. The Later Mesolithic Population of Sjaelland, Denmark, and the Neolithic Transition. In: Svejle M, Domanska L, and Dennell R, editors. *Harvesting the Sea, Farming the Forest: The Emergence of Neolithic Societies in the Baltic Region*. Sheffield: Sheffield Academic Press. p 203-212.
- Merzoug S. 2014. A level prior to the Upper Capsian at Medjez II (Algeria): Archaeozoological and taphonomical evidence combined with archaeological data. *Quaternary International* 320(0):125-130.
- Morales J. (in this volume) The contribution of plant macro-remains to the study of the Palaeolithic diet: an insight from the Later Stone Age of North West Africa. *Journal of Archaeological Science: reports*.
- Mulazzani S, Belhouchet L, Salanova L, Aouadi N, Dridi Y, Eddargach W, Morales J, Tombret O, Zazzo A, and Zoughlami J. 2016. The emergence of the Neolithic in North Africa: A new model for the Eastern Maghreb. *Quaternary International*. 410: 123-143.
- Nelson GC. 2015. A Host of Other Dental Diseases and Disorders. *A Companion to Dental Anthropology*: John Wiley & Sons, Inc. p 465-483.
- Ortner DJ, and Putschar W. 1981. *Identification of Pathological Conditions in Human Skeletal Remains*. Washington D.C.: Smithsonian Institution Press. 488 p.
- Oyamada J, Kitagawa Y, Manabe Y, and Rokutanda A. 2004. Dental pathology in the samurai and commoners of early modern Japan. *Anthropological Science* 112(3):235-246.
- Pinhasi R, Eshed V and Shaw P. 2008. Evolutionary changes in the masticatory complex following the transition to farming in the southern Levant. *American Journal of Physical Anthropology*, 135(2): 136-148.
- Pomeroy E and Zakrzewski SR. 2009. Sexual dimorphism in diaphyseal cross-sectional shape in the medieval Muslim population of Écija, Spain, and Anglo-Saxon Great Chesterford, UK. *International Journal of Osteoarchaeology* 19(1): 50-65.
- Rahmani N. 2004. Technological and cultural change among the last hunter-gatherers of the Maghreb: the Capsian (10,000–6000 BP). *Journal of World Prehistory* 18(1):57-105.
- Roche J. 1963. *L'Épipaléolithique Marocain*: Fondation Calouste Gulbenkian. 261 p.
- Royer P. 1926. Crâne préhistorique de la grotte des hyènes (Constantine). *Bulletins et Mémoires de la Société d'anthropologie de Paris* 7(4):119-120.
- Ruff, C. B., Larsen, C. S. and Hayes, W. C. (1984) 'Structural changes in the femur with the transition to agriculture on the Georgia coast', *American Journal of Physical Anthropology*, 64(2), 125-136.

- Schollmeyer KG, and Turner C. 2004. Dental Caries, Prehistoric Diet, and the Pithouse-to-Pueblo Transition in Southwestern Colorado. *American Antiquity* 69(3):569-582.
- Schwenninger JL, Collcut SN, Barton RNE, Bouzouggar A, El Hajraoui A, Nespoulet R, Debenath A, and Clark-Balzan L. 2010. Luminescence chronology for Aterian cave sites on the Atlantic coast of Morocco. In: Garcea EEA, editor. *South-Eastern Mediterranean Peoples between 130,000 and 10,000 Years Ago*. Oxford: Oxbow Books. p 18-36.
- Sheppard P, and Lubell D. 1991. Early Holocene Maghreb prehistory: an evolutionary approach. *Sahara* 3:63-69.
- Starling, A. P. and Stock, J. T. (2007) 'Dental indicators of health and stress in early Egyptian and Nubian agriculturalists: A difficult transition and gradual recovery', *American Journal of Physical Anthropology*, 134(4), 520-528.
- Tayles N. 2009. Can dental caries be interpreted as evidence of farming? The Asian experience. *Frontiers of Oral Biology* 13:162-166.
- Tayles N, Domett K, and Nelsen K. 2000. Agriculture and dental caries? The case of rice in prehistoric Southeast Asia. *World Archaeology* 32(1):68-83.
- Taylor V, Barton R, Bell M, Bouzouggar A, Collcutt S, Black S, and Hogue J. 2011. The Epipalaeolithic (Iberomaurusian) at Grotte des Pigeons (Taforalt), Morocco: a preliminary study of the land Mollusca. *Quaternary International* 244(1):5-14.
- Temple DH. 2015. Caries. *A Companion to Dental Anthropology*: John Wiley & Sons, Inc. p 433-449.
- Temple DH, and Larsen CS. 2007. Dental caries prevalence as evidence for agriculture and subsistence variation during the Yayoi period in prehistoric Japan: Biocultural interpretations of an economy in transition. *American Journal of Physical Anthropology* 134(4):501-512.
- Turner CG. 1979. Dental anthropological indications of agriculture among the Jomon people of central Japan. X. Peopling of the Pacific. *American Journal of Physical Anthropology* 51(4):619-635.
- Turner CG, and Machado LMC. 1983. A new dental wear pattern and evidence for high carbohydrate consumption in a Brazilian Archaic skeletal population. *American Journal of Physical Anthropology* 61(1):125-130.
- Vallois HV and Felice S. 1979. Le squelette Capsien d'Ain Meterchem (Tunisie). *L'Anthropologie* 83: 395-416
- Watson JT 2008. Prehistoric dental disease and the dietary shift from cactus to cultigens in northwest Mexico. *International Journal of Osteoarchaeology* 18(2):202-212.

List of Figures and Tables in the Manuscript

Figure 1. Map showing site locations included in the sample.


Figure 2. Frequency of carious lesions (per total number of teeth in past populations) grouped according to subsistence pattern: Hunting and Gathering; Mixed Economy including horticulture, agriculture + hunting, gathering or fishing); and agriculture (food producing) compared to the Iberomaurusians and Capsians (Adapted from Lanfranco and Eggers 2010).

Table 1. Sample composition.

Table 2. Frequencies of oral pathologies by tooth sockets and observed teeth.

Table 3. Frequencies of oral pathologies by individual.

Table 4. Sample size and caries frequencies in two Iberomaurusian samples: Taforalt and Afalou.






Table 1: Sample composition

Iberomaurusian	Iberomaurusian continued	Iberomaurusian continued
Age category	Age category	Age category
<i>Young Adult</i>	<i>Old Adult</i>	<i>Undetermined continued (UD)</i>
Afalou 1	Afalou 10	Taforalt XX
Afalou 20	Afalou 27	Taforalt XXI-1
Afalou 22	Afalou 40	Taforalt XXIV-2
Afalou 23	Afalou 49	Taforalt XXV-C3-2+C3
Afalou 24	Taforalt IX-1	Taforalt XXVI
Afalou 29	Taforalt XV-C2	Taforalt XXVII-C1
Afalou 31	Taforalt XXV-C1	Taforalt-no number
Afalou 38	<i>Undetermined (UD)</i>	
Afalou 48	Afalou 1935 Md 6 R	Capsian
Taforalt Ind 5	Afalou 1936 - 6 "0"	Age category
Taforalt VIII-1	Afalou 1936 c	<i>Young Adult</i>
Taforalt XVII-C1-C3	Afalou 34	Ain Dokkhara 1
Taforalt XVII-C3 (max)	Afalou 42	<i>Adult</i>
Taforalt XX-1	Afalou 7	Khanguet al Mouhaad IV
Taforalt XXV C4 + SXXV	Afalou 8 s.m.	Ain Meterchem 1
<i>Adult</i>	Afalou A5ho	Site 12 3A1
Afalou 11	Afalou As	Site 12 3A-2
Afalou 12	Afalou B4	Site12 3A-5
Afalou 13	Afalou C Md8	<i>Old Adult</i>
Afalou 14	Afalou D6 Ma Fem	Hyenes 4-255 mand
Afalou 15	Afalou F Md10	Mechta type male
Afalou 17	Afalou I 1935-6	<i>Undetermined (UD)</i>
Afalou 2	Afalou Max 1935-6a	Hyenes 4-254 skull
Afalou 25	Afalou Max 1935b	Hyenes 4-255
Afalou 28	Afalou max 1936-d	Hyenes D
Afalou 3	Afalou Maxilla 35	Hyenes G
Afalou 30	Afalou Md 2-3	Koudiat-el-Kherrouba
Afalou 32	Afalou Md4b	Mechta 4
Afalou 36	Afalou Md5	Mechta 5
Afalou 39	Afalou Mx4	Mechta max mand part
Afalou 43	Dar Es Soltan I	Mesloug
Afalou 45	Taforalt Ind 1	Redeyef A
Afalou 46	Taforalt Ind 10	Site 12 Mand frag
Afalou 47	Taf iso mand	
Afalou 5	Taforalt I 1952	
Afalou 9	Taforalt I- no number (a)	
Taforalt Ind 13	Taforalt I- no number (b)	
Taforalt Ind 14	Taforalt innomine	
Taforalt Ind 3	Taforalt IX-B	
Taforalt Ib	Taforalt VIII ?	
Taforalt trepanated	Taforalt XI-B-C2	
Taforalt VIII-2	Taforalt XII-?	
Taforalt XI-1	Taforalt XIV-1	
Taforalt XII-1	Taforalt XIX	
Taforalt XII-C4	Taforalt XIX-C1	
Taforalt XV-C4C5	Taforalt XIX-C2	
Taforalt XVIII-1	Taforalt XV-C5	
Taforalt XX-C1	Taforalt XV-C5max	
Taforalt XXII-1	Taforalt XVI C1 NOPD mand	
Taforalt XXIV-1	Taforalt XVI C1 NOPD max	
Taforalt XXV-C3.1 + C3	Taforalt XVI-C2	
Taforalt XXVII-1	Taforalt XVII-C2	

Table 2: Frequencies of oral pathologies by tooth sockets and observed teeth.

	No. Of observable sockets		No. Of observable PC sockets *		No. sockets with PMTL*		No. of valid observable Sockets*		No. of PC sockets with AMTL~		No. of observed teeth*		No. teeth with caries^		No. teeth with gross caries\$		No. teeth with crown caries\$		No. teeth with root caries\$		No. of teeth with resorption^		No. of sockets with abscess*	
	n		n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%
Iberomaurusian																								
YA	350		222	63.4	49	14.0	301	86.0	7	3.2	273	78.0	115	42.1	4	3.5	64	55.7	14	12.2	136	49.8	6	1.7
MA	752		468	62.2	89	11.8	663	88.2	49	10.5	557	74.1	364	65.4	14	3.8	104	28.6	33	31.7	320	57.5	32	4.3
OA	176		113	64.2	17	9.7	159	90.3	24	21.2	124	70.5	84	67.7	10	11.9	11	13.1	6	54.5	55	44.4	21	11.9
UA	713		455	63.8	179	25.1	534	74.9	32	7.0	463	64.9	281	60.7	13	4.6	93	33.1	49	0.0	216	46.7	28	3.9
Total	1991		1258	63.2	334	16.8	1657	83.2	112	8.9	1417	71.2	844	59.6	41	4.9	272	32.2	102	37.5	727	51.3	87	4.4
Capsian																								
YA	32		20	62.5	1	3.1	31	96.9	0	0.0	31	96.9	22	71.0	0	0.0	7	31.8	0	0.0	32	103.2	1	3.1
MA	142		88	62.0	6	4.2	136	95.8	13	14.8	103	72.5	39	37.9	3	7.7	15	38.5	9	60.0	56	54.4	14	9.9
OA	35		23	65.7	6	17.1	29	82.9	4	17.4	21	60.0	11	52.4	1	9.1	7	63.6	3	42.9	17	81.0	2	5.7
UA	195		119	61.0	34	17.4	161	82.6	17	14.3	116	59.5	61	52.6	0	0.0	14	23.0	3	21.4	53	45.7	13	6.7
Total	404		250	61.9	47	11.6	357	88.4	34	13.6	271	67.1	133	49.1	4	3.0	43	32.3	15	34.9	158	58.3	30	7.4

Young adults (YA), middle adults (MA), old adults (OA), Undetermined age (UA)

* % of no. of observable sockets

% of no. of valid observed sockets (=observed sockets - PMTL)

~ % of no. observable PC sockets

^ % of no. of observed teeth

\$ % of no. of teeth with caries

Table 3: Frequencies of oral pathologies by individual.

	No. of ind.	No. of ind. with PC sites §		No. of ind. with PMTL §		No. of ind. with PC AMTL sites §		No. of ind. with resorption §		No. of ind. with abscess §		No. of ind. with caries §		No. of ind. with gross caries #	
Iberomaurusian	n	n	%	n	%	n	%	n	%	n	%	n	%	n	%
YA	15	15	100.0	10	66.7	4	31.0	12	80.0	6	40.0	14	93.3	2	14.3
MA	36	36	100.0	20	55.6	16	44.0	34	94.4	15	41.7	36	100.0	9	25.0
OA	7	7	100.0	5	71.4	6	86.0	6	85.7	5	71.4	7	100.0	2	28.6
UA	51	51	100.0	44	86.3	44	86.0	41	80.4	14	27.5	49	96.1	10	20.4
Total	109	109	100.0	78	71.6	38	35.0	93	85.3	40	36.7	106	97.2	23	21.7
Capsian															
YA	1	1	100.0	1	100.0	0	0.0	1	100.0	1	100.0	1	100.0	0	0.0
MA	5	5	100.0	4	80.0	4	80.0	3	60.0	4	80.0	4	80.0	2	50.0
OA	2	2	100.0	2	100.0	1	50.0	2	100.0	1	50.0	1	50.0	1	100.0
UA	11	11	100.0	9	81.8	5	45.5	7	63.6	6	54.5	8	72.7	0	0.0
Total	19	19	100.0	16	84.2	10	52.6	13	68.4	12	63.2	14	73.7	3	21.4

§ % of no. of individuals

% of ind with teeth with caries

Table 4: Sample size and caries frequencies in two Iberomaurusian samples: Taforalt and Afalou.

	No. Of observable sockets	No. sockets with PMTL	No. sockets with AMTL	No. of observed teeth	No. teeth with caries^	
Afalou	n	n	n	n	n	%
YA	183	24	18	141	71	50.4
MA	465	53	83	329	239	72.6
OA	90	8	13	69	55	79.7
UA	286	68	30	188	133	70.7
Total	1024	153	144	727	498	68.5
Taforalt						
YA	167	25	10	132	44	33.3
MA	287	36	23	228	125	54.8
OA	86	9	22	55	29	52.7
UA	411	105	40	266	147	55.3
Total	951	175	95	681	345	50.7

Young adults (YA), middle adults (MA), old adults (OA), Undetermined age (UA)

^ % of no. of observed teeth (observed teeth = observable sockets - AMTL - PMTL)